

Muon anomalous magnetic moment constrains models with excited leptons

Subhendu Rakshit *

*Harish-Chandra Research Institute, Chhatnag Road, Jhusi,
Allahabad 211 019, India*

Abstract

We explore the possibility of the existence of excited leptons in the light of recent data on muon $g - 2$ from BNL. We have been able to put stringent bounds on the relevant parameter space.

1 Introduction

The recent measurement of the anomalous magnetic moment (a_μ) of the muon by the E821 experiment [1] at BNL seem to indicate a 2.6σ deviation from theoretical predictions based on the Standard Model (SM), more precisely, $\delta a_\mu \equiv a_\mu^{expt} - a_\mu^{SM} = 426(165) \times 10^{-11}$. The ongoing analysis of collected data and accumulation of fresh data is likely to reduce the errors even further. The required additional contribution to explain this anomaly is nearly three times the standard electroweak contribution. This deviation might be indicative of some new physics beyond the Standard Model (SM) and this has led to consideration of several extensions of the SM [2]. Of course one should keep in mind that the existing SM calculations for the hadronic contributions to a_μ are questionable [3] and there is no universal consensus amongst different groups performing these calculations. But this does not undermine the credibility of exploring new physics options beyond the SM.

In the SM leptons and quarks are treated as fundamental point-like objects. But ordinary leptons might be made up of some more fundamental particles [4] interacting in a very strong confining interaction characterized by some scale $\Lambda \sim \mathcal{O}(1)$ TeV. A possible prediction of these composite models [5] is the existence of excited states of ordinary leptons. It was pointed out in Ref. [6] that these excited leptons can give rise to a large magnetic moment for the muon. The reported anomaly on a_μ can be explained in the presence of excited leptons and this, in turn, severely restricts the allowed parameter space.

*E-mail address: srakshit@mri.ernet.in

2 Description of the model

We consider excited state of the muon (μ^*) and its corresponding excited neutrino (ν_μ^*) to contribute at one loop to a_μ [7]. In this study we shall confine ourselves to the lowest lying excited states of spin and isospin 1/2, although in principle, other excited states with higher spin and various isospin values might also exist [8]. However, this is sufficient for a *conservative order of magnitude estimation* as we do not see any compelling reason for any fine cancellation between the contributions from the first excited state and other excited states.

These excited muons are very massive in comparison to the ordinary ones. To motivate this large mass gap, we assume that these excited states acquire their masses prior to $SU(2) \times U(1)$ breaking and so both of their left- and right-handed states will be in weak isodoublets. As a consequence, the interactions of μ^* and ν_μ^* with the gauge bosons are vector-like [9]. The corresponding Lagrangian can be written as,

$$\mathcal{L}_{VF'F} = - \sum_{V=\gamma, Z, W} C_{VF'F} \bar{F}' \gamma_\mu F V^\mu + \text{h.c.}, \quad (1)$$

where $F, F' = \mu^*, \nu_\mu^*$. The constants $C_{VF'F}$'s are given by

$$\begin{aligned} C_{Z\mu^*\mu^*} &= \frac{g}{2 \cos \theta_W} (1 - 2 \sin^2 \theta_W), & C_{Z\nu_\mu^*\nu_\mu^*} &= -\frac{g}{2 \cos \theta_W} \\ C_{\gamma\mu^*\mu^*} &= e & C_{W\nu_\mu^*\mu^*} &= -\frac{g}{\sqrt{2}} \end{aligned}$$

In the above Lagrangian and in the following ones, we will indicate only those interactions relevant for our subsequent discussions.

On the other hand, the Lagrangian describing the transition between the ordinary muon and the excited muon may look like:

$$\begin{aligned} \mathcal{L}_{Ff} &= \frac{1}{4\Lambda} \bar{\mathbf{F}}_{\mathbf{R}} \sigma^{\mu\nu} [a \boldsymbol{\tau} \cdot \mathbf{W}_{\mu\nu} + b Y B_{\mu\nu}] \mathbf{f}_{\mathbf{L}} \\ &+ \frac{1}{4\Lambda^2} \bar{\mathbf{F}}_{\mathbf{L}} \sigma^{\mu\nu} [a' \boldsymbol{\tau} \cdot \mathbf{W}_{\mu\nu} + b' Y B_{\mu\nu}] f_R \Phi + \text{h.c.} . \end{aligned} \quad (2)$$

It is customary to parametrize the new coupling parameters as $a = g f_L$, $a' = g' f'_L$, $b = g f_R$ and $b' = g' f'_R$. Here, $\{f_L, f_R\}$ and $\{f'_L, f'_R\}$ are the weight factors associated with the gauge groups $SU(2)$ and $U(1)$ respectively and they arise from the underlying dynamics describing the compositeness. However, in this paper, for the sake of convenience, we shall be using the coupling parameters a, b, a' and b' instead of f_L, f_R, f'_L and f'_R .

In the above Lagrangian,

$$\mathbf{F}_{\mathbf{L}} = \begin{pmatrix} \nu_\mu^* \\ \mu^* \end{pmatrix}_L, \quad \mathbf{F}_{\mathbf{R}} = \begin{pmatrix} \nu_\mu^* \\ \mu^* \end{pmatrix}_R, \quad \mathbf{f}_{\mathbf{L}} = \begin{pmatrix} \nu_\mu \\ \mu \end{pmatrix}_L, \quad f_R = \mu_R.$$

Φ is the SM Higgs doublet.

It is important to note in $\mathcal{L}_{Ff}$ that we have included both left- and right-handed excited leptons. It was observed in Ref. [6] that this might lead to a large magnetic

moment for the ordinary leptons. To avoid this, it is a common practice to consider only the right-handed excited fermions interacting this way. Here we allow both type of couplings to review the bounds imposed on the coupling parameters in view of the recent anomaly in the magnetic moment of the muon. Inclusion of left-handed excited leptons require consideration of operators of higher dimensionality than those for right-handed ones. So the interactions arising from second line of eqn. 2 will suffer an additional suppression factor of v/Λ in comparison to those from the first line. Here v stands for the VEV of the SM Higgs.

It is also possible to include a term like

$$\frac{1}{4\Lambda^2} \bar{\mathbf{F}}_{\mathbf{L}} \sigma^{\mu\nu} [a'' \boldsymbol{\tau} \cdot \mathbf{W}_{\mu\nu} + b'' Y B_{\mu\nu}] \nu_R \tilde{\Phi} + \text{h.c.} \quad (3)$$

in eqn. 2, which needs introduction of two more couplings a'' and b'' . For the sake of minimality, we set them to zero. In the above $\tilde{\Phi} \equiv -i\sigma_2\Phi^*$, σ_2 being the second Pauli matrix.

The interaction Lagrangian for μ and ν_μ^* with a single gauge boson obtained from eqn. 2 looks like:

$$\mathcal{L}_{VFf} = -\frac{1}{2\Lambda} \sum_{V=\gamma, Z, W} \bar{F} \sigma_{\mu\nu} (D_{VFf}^L P_L + D_{VFf}^R P_R) f \partial^\mu V^\nu + \text{h.c.}, \quad (4)$$

where $f = \mu, \nu_\mu$. The constants D_{VFf} 's are given by,

$$\begin{aligned} D_{Z\mu^*\mu}^L &= -a \cos \theta_W + b \sin \theta_W & D_{Z\mu^*\mu}^R &= \frac{v}{\Lambda} (-a' \cos \theta_W + b' \sin \theta_W) \\ D_{\gamma\nu_\mu^*\nu_\mu}^L &= a \sin \theta_W - b \cos \theta_W & D_{\gamma\nu_\mu^*\nu_\mu}^R &= 0 \\ D_{\gamma\mu^*\mu}^L &= -a \sin \theta_W - b \cos \theta_W & D_{\gamma\mu^*\mu}^R &= \frac{v}{\Lambda} (-a' \sin \theta_W - b' \cos \theta_W) \\ D_{W\nu_\mu^*\mu}^L &= D_{W\mu^*\nu_\mu}^L = a\sqrt{2} & D_{W\nu_\mu^*\mu}^R &= \frac{v}{\Lambda} a\sqrt{2}; \quad D_{W\mu^*\nu_\mu}^R = 0 \end{aligned}$$

The Lagrangian describing quartic interactions between μ and μ^* with W and γ arising from the non-abelian part of $W_{\mu\nu}$ in eqn. 2 can be written as,

$$\mathcal{L}_{VVFf} = -\frac{1}{2\Lambda} \bar{F} \sigma_{\mu\nu} (E_{VVFf}^L P_L + E_{VVFf}^R P_R) f W^\nu + \text{h.c.}, \quad (5)$$

where the constants E_{VVFf} 's are given by

$$E_{\gamma W\mu^*\nu_\mu}^L = -E_{\gamma W\nu_\mu^*\mu}^L = -a e\sqrt{2}; \quad E_{\gamma W\mu^*\nu_\mu}^R = 0; \quad E_{\gamma W\nu_\mu^*\mu}^R = \frac{v}{\Lambda} a' e\sqrt{2}$$

For the sake of notational convenience, we also define the required SM interactions in the following Lagrangian:

$$\mathcal{L}_{Vf'f} = - \sum_{V=\gamma, Z, W} \bar{f}' \gamma^\mu (F_{Vf'f}^L P_L + F_{Vf'f}^R P_R) f. \quad (6)$$

3 Excited lepton contribution to a_μ

With the interactions defined in the previous section, we can calculate the one-loop diagrams containing excited leptons contributing to a_μ . It has to be pointed out that in the SM, the self-energy type diagrams do not contribute to a_μ but here diagrams 5—8 (see Fig. 1) contribute due to the transition magnetic interactions between ordinary and excited leptons. It is obvious from the Dirac structure of the vertices that the longitudinal parts of the gauge bosons do not contribute. As a consequence, in the absence of form-factors (see the following discussion), one has to encounter at most quadratic divergences.

There are several alternatives to make the momentum integrals convergent. One of the options is to apply a momentum cut-off at $k^2 = \Lambda^2$ [6]. Another option is to use form-factors [10], as one can no longer treat the leptons as point particles. We shall follow this alternative. We associate dipole form-factors $\left(\frac{1}{1-k^2/\Lambda^2}\right)^2$ with the vertices which involve magnetic transitions between the excited leptons with the ordinary ones. Here k denotes the momentum of the associated off-shell gauge boson. Muon magnetic moment calculation in the dimensional regularization method was performed for excited leptons (with a different set of interactions) in Ref. [11]. We make a remark in passing that for an order of magnitude estimation, the numerical differences coming from the choice of different regularization schemes are not important [6].

In the appendix we present analytical expressions of the contributions to a_μ from the dominant diagram sets. The contributions from the diagrams 9 and 10 containing the quartic coupling are always insignificant. Contribution from diagrams 5—8 can be more than 90% of the total excited lepton contribution to a_μ . When a and a' both are zero, the W diagrams do not contribute as the $SU(2)$ sector does not participate in the transition magnetic interaction between the ordinary and the excited leptons. From the symmetry of the diagrams it is clear that diagram sets 3 and 4 give equal contributions. The same is true for the other sets $\{5, 6\}$, $\{7, 8\}$ and $\{9, 10\}$.

In this model there are several extra parameters than the standard model: (a) the coupling parameters a , b , a' and b' , (b) the masses of μ^* and ν_μ^* and (c) the compositeness scale — Λ . We take μ^* and ν_μ^* to be degenerate in mass (M_F), so that they do not contribute to the ρ parameter.

4 Results and discussion

In this section we shall mainly divide our attention onto two categories of couplings. At first we shall deal with the situation when only one of the coupling constants is nonzero. In this situation the transition magnetic interactions between the ordinary and the excited leptons will respect a chiral symmetry, *i.e.*, only left- or the right-handed excited lepton will be coupled to ordinary fermions. We shall refer to this as chirality conserving interactions. The other obviously will be chirality violating and will arise when either a and a' or b and b' will be simultaneously non-zero. Magnetic moment being a chirality violating




Figure 1: *Diagram sets containing excited leptons contributing to a_μ*

operator one need to have a left-handed muon at one end of the Feynman diagram and a right-handed muon on the other. Now in each diagram there are two transition magnetic interactions. So to have a muon magnetic moment one needs to have at least one mass insertion on the fermion line. For the chirality violating interactions, one can have a mass insertion on the excited lepton line making the contribution large. But for the chirality conserving case, one gets a helicity flip on the muon line making the contribution more suppressed. We can now proceed to discuss the parameter dependence of δa_μ .

4.1 Dependence on coupling constants

Now let us discuss the parameter dependence of the total contribution to a_μ from these diagrams. If we consider only a to be nonzero, then a_μ increases quadratically with it, *i.e.*, $\delta a_\mu = y(a/\Lambda)^2$. For $M_F = 400$ GeV and $\Lambda = 1$ TeV, $y(\text{in GeV}^2) = 1.6 \cdot 10^{-3}$. For b , a' and b' , y is evaluated as $5.8 \cdot 10^{-4}$, $4.3 \cdot 10^{-5}$ and $5.5 \cdot 10^{-5}$ respectively.

For chirality violating case, it turns out that the terms containing the products aa' or bb' give the dominant contribution due to a mass insertion on the excited lepton line. Now for $a = a'v/\Lambda$ or $b = b'v/\Lambda$, the magnetic transition between the ordinary and excited leptons are pure vectorial, whereas for $a = -a'v/\Lambda$ or $b = -b'v/\Lambda$, the interactions are axial-vector-like. We shall deal with these two extremes and in all the above cases the dependence of δa_μ on a or b is obviously quadratic as before. $y = 9.9$ for $a = a'v/\Lambda$ and $y = 5.3$ for $b = b'v/\Lambda$. Comparing these values of y with the previous case it is clear that in this case there will be a large contribution from excited leptons.

4.2 Dependence on M_F

For chirality conserving case, δa_μ is almost proportional to $1/M_F^2$ (see Fig. 2(a)). The reason behind is that the diagram sets 5—8 give the dominant contributions. In these diagrams, we can pick $1/M_F^2$ from the propagator joining the one-loop part to the external photon vertex. As discussed earlier, one can have a helicity flip on the muon line only. So no other factor of M_F comes into the picture. M_F dependence coming from the propagators inside the loops makes a little deviation from $1/M_F^2$. But for chirality violating case, there is another power of M_F in the numerator due to a helicity flip. Hence, then the dependence is more like $1/M_F$ (see Fig. 2(b)).

4.3 Dependence on Λ

In Fig. 3, we explore the dependence of δa_μ on the compositeness scale Λ . In fig. (a) we see that the contribution increases and then rather than blowing up saturates as we keep on increasing Λ . Here we have kept M_F fixed at 400 GeV. But if we take $M_F \sim \Lambda$, then it falls rapidly (see fig. (b)). The reason is that in this case the M_F dependence also adds up to the aforesaid Λ dependence. With respect to this situation, the plotted behaviour in (a) is rather flat.




Figure 2: Dependence of a_μ on excited lepton mass. In (a) we have taken either a or b to be non-zero. In (b) we have considered the chirality violating cases when $a = a'v/\Lambda$ or $b = b'v/\Lambda$. We have put $\Lambda = 1$ TeV.




Figure 3: Dependence of δa_μ on the compositeness scale Λ . In diagram (a) we have kept M_F fixed at 400 GeV. In (b) we take $M_F \sim \Lambda$.

4.4 Constraints on the parameter space

To get an idea of the parameter space allowed by the recent measurement on a_μ , it is convenient to look at the contour plots for δa_μ in the coupling constant — M_F plane. Let us discuss the situation when only one of the couplings is non-zero. For simplicity of presentation, we take only a to be non-zero. Now we have seen that in this case $\delta a_\mu \propto a^2$. The M_F dependence is more like $1/M_F^2$. So we expect almost linear behaviour in the contours $a \sim z_1 \sqrt{\delta a_\mu} M_F$. z_1 (in GeV^{-1}) is evaluated (for $\Lambda = 1 \text{ TeV}$) as 66, 108, 450 and 342 for a, b, a' and b' respectively. For an order of magnitude estimation this formula works quite well. In Fig. 4(a), we indicate regions in the $a - M_F$ plane allowed by the recent data on $g - 2$ measurement with their corresponding confidence levels.




Figure 4: Contour plots for fixed values of δa_μ in M_F Vs. coupling constant plane.

Now let us deal with two non-zero coupling constants. We shall start with the situation when the excited leptons have pure vectorial transition magnetic coupling with the ordinary leptons. In this case the contribution will be positive. Argument in the same line as before leads to a relation $a \sim z_2 \sqrt{\delta a_\mu} M_F$. z_2 (in $\text{GeV}^{-1/2}$) is given by 16.5 and 22.9 for $a = a'v/\Lambda$ and $b = b'v/\Lambda$ respectively.

In Fig. 4(b), we present the contour plot for $a = a'v/\Lambda$. We see that the constraints arising out in this situation are more stringent. If one of them is negative, the total contribution is also negative which is allowed at the most at a level of 3σ .

5 Summary and conclusions

We have discussed both types of transition magnetic moment-type coupling of excited leptons to ordinary leptons – (a) chirality conserving and (b) chirality violating. For the later type of coupling, one gets a huge contribution to a_μ [6]. We have explored how these models with excited leptons can explain the recent anomaly on muon anomalous magnetic

moment measurement. This puts stringent constraints on the relevant parameter space.

Excited leptons can also contribute radiatively in the muon mass. Any such correction to the muon mass can be absorbed in the bare muon mass using on-shell renormalization conditions. In the chirality violating case (which is not a natural theory), one can radiatively generate muon mass this way even taking tree level muon mass to be zero. There the contribution will be large due to a factor of M_F coming from a helicity flip on the muon line and it will not be possible to absorb it in the bare mass [12], which is zero. Consequently, one can get stringent bounds on the coupling parameters for a given compositeness scale $\Lambda \sim \mathcal{O}(1)$ TeV. However, here we do not try to generate muon mass in this way. In the chirality violating case, the weight factors — f_i 's turn out to be very small and this has to be ensured by the underlying dynamics describing compositeness.

We find that for the chirality violating scenario, the values of the parameters a and a' (or b and b') with a relative negative sign are disfavoured by the data. The constraints we get are more stringent than the limits obtained from direct searches of excited leptons at LEP, where for unit transition magnetic moment-type couplings, the lower bound on excited lepton mass is $\mathcal{O}(100)$ GeV [13]. This rules out any hope for finding these particles via direct searches, if they have chirality violating couplings. On the other hand, the scenario for particles having chirality conserving couplings may not be that grim, but the recent BNL measurement of a_μ leaves behind a very small allowed parameter space. Recently search prospects for excited leptons have been studied for the Fermilab Tevatron [14] and CERN Large Hadron Collider [15]. The studies have been made for e^* and ν_e^* in the chirality conserving scenario. However, if we take the electron and muon detection efficiencies to be similar then the same bounds can be realized for excited muons as well. In the case of the Tevatron (run II), it has been claimed that for $f_L/\Lambda > 1$ TeV⁻¹, an excited lepton mass less than 250 GeV can be ruled out at 95% CL. For LHC, the bounds will be far more stringent. It will be capable of ruling out a region (at 95% CL) $f_L/\Lambda > 0.7$ TeV⁻¹ for excited lepton masses less than 1 TeV and will be sensitive upto 2 TeV. Hence, we see that if excited leptons are the culprit for the recent muon anomaly, they can leave their footprints in the Tevatron or in the LHC, if they have chirality conserving transition magnetic coupling with the ordinary leptons. But even if these colliders fail to detect them, the excited leptons can still explain the muon anomaly if they have small chirality violating couplings.

Acknowledgements

S.R. would like to thank Debajyoti Choudhury for suggesting this work, help in algebraic computation and for numerous comments and discussions. He thanks Biswarup Mukhopadhyaya, Uma Mahanta, Debrupa Chakraverty and Anindya Datta for illuminating discussions. He also thanks the Abdus Salam International Centre for Theoretical Physics, Italy for hospitality where a major part of this work was carried out.

Appendix

The total contribution of excited leptons to anomalous magnetic moment of muon is given by,

$$\delta a_\mu = \frac{m_\mu}{16\pi^2} \Lambda^2 \left[\int_0^1 dx_1 \int_0^{1-x_1} dx_2 \int_0^{1-x_1-x_2} dx_3 \sum_{i=1}^4 X_i + \int_0^1 dx_1 \int_0^{1-x_1} dx_2 \sum_{i=5}^8 X_i \right]$$

The X_i 's are contributions from different diagram sets (Fig. 1).

$$\begin{aligned} X_1 &= 8\Lambda^4 x_1^3 \left(\frac{f_{1b} M_F (1 - 9x_1 - 9x_2)}{\Delta_1^4} \right. \\ &\quad \left. - m_\mu (1 - x_1 - x_2) f_{1a} \left(-\frac{8M_F^2 (1 - x_1 - x_2)}{\Delta_1^5} + \frac{(5 + 4x_1 + 4x_2)}{\Delta_1^4} \right) \right) \\ X_2 &= -8\Lambda^4 (1 - x_1 - x_2 - x_3)^3 \frac{(3f_{2b} M_F (3x_2 - 2) - f_{2a} m_\mu x_2 (13 - 12x_2))}{\Delta_2^4} \\ X_3 &= 24x_1 \left(-\frac{2f_{3c} M_F}{3\Delta_3^2} - \frac{2f_{3d} m_f M_F m_\mu}{3\Delta_3^3} (1 - x_1 - x_3) + \frac{f_{3a} m_\mu}{\Delta_3^2} (x_1 + x_3) \right) \\ X_4 &= 24x_1 \left(-\frac{2f_{4b} M_F}{3\Delta_4^2} - \frac{2f_{4d} m_f M_F m_\mu}{3\Delta_4^3} (1 - x_1 - x_3) + \frac{f_{4a} m_\mu}{\Delta_4^2} (x_1 + x_3) \right) \\ X_5 &= 24x_1 \left(\frac{2f_{5b} M_F}{\Delta_5} + \frac{2f_{5a} m_\mu}{\Delta_5} + \frac{f_{5a} M_F M_{F'} m_\mu x_2}{\Delta_5^2} \right) \\ X_6 &= 24x_1 \left(\frac{2f_{6b} M_F}{\Delta_6} + \frac{2f_{6a} m_\mu}{\Delta_6} + \frac{f_{6a} M_F M_{F'} m_\mu x_2}{\Delta_6^2} \right) \\ X_7 &= 24x_1 \left(\frac{2f_{7b} M_F}{\Delta_7} + \frac{2f_{7a} m_\mu}{\Delta_7} + \frac{f_{7d} m_f M_F m_\mu x_2}{\Delta_7^2} \right) \\ X_8 &= 24x_1 \left(\frac{2f_{8c} M_F}{\Delta_8} + \frac{2f_{8a} m_\mu}{\Delta_8} + \frac{f_{8d} m_f M_F m_\mu x_2}{\Delta_8^2} \right) \end{aligned}$$

The Δ_i 's are given by,

$$\begin{aligned} \Delta_1 &= -m_V^2 x_2 - M_F^2 (x_1 + x_3) - \Lambda^2 (1 - x_1 - x_2 - x_3) \\ \Delta_2 &= -M_F^2 x_2 - m_V^2 (x_1 + x_3) - \Lambda^2 (1 - x_1 - x_2 - x_3) \\ \Delta_3 &= -\Lambda^2 x_1 - m_V^2 x_3 - M_F^2 (1 - x_1 - x_2 - x_3) \\ \Delta_4 &= -\Lambda^2 x_1 - M_F^2 x_2 - m_V^2 x_3 \\ \Delta_5 &= \Delta_6 = -\Lambda^2 x_1 - M_{F'}^2 x_2 - m_V^2 (1 - x_1 - x_2) \\ \Delta_7 &= \Delta_8 = -\Lambda^2 x_1 - m_V^2 (1 - x_1 - x_2) \end{aligned}$$

The f 's are defined as

$$\begin{aligned}
f_{1a} &= \frac{1}{32} ((D_{V\mu^*\mu}^L)^2 + (D_{V\mu^*\mu}^R)^2) & f_{1b} &= \frac{1}{16} D_{V\mu^*\mu}^L D_{V\mu^*\mu}^R \\
f_{2a} &= -\frac{1}{32} ((D_{V\nu_\mu^*\mu}^L)^2 + (D_{V\nu_\mu^*\mu}^R)^2) & f_{2b} &= \frac{1}{16} D_{V\nu_\mu^*\mu}^L D_{V\nu_\mu^*\mu}^R \\
f_{3a} &= \frac{1}{32e} (D_{VF\mu}^L D_{\gamma Ff}^L F_{Vf\mu}^L + L \leftrightarrow R) & f_{3b} &= \frac{1}{32e} (D_{VF\mu}^L D_{\gamma Ff}^L F_{Vf\mu}^R + L \leftrightarrow R) \\
f_{3c} &= \frac{1}{32e} (D_{VF\mu}^L D_{\gamma Ff}^R F_{Vf\mu}^R + L \leftrightarrow R) & f_{3d} &= \frac{1}{32e} (D_{VF\mu}^L D_{\gamma Ff}^R F_{Vf\mu}^L + L \leftrightarrow R) \\
f_{4a} &= \frac{1}{32e} (F_{Vf\mu}^R D_{\gamma Ff}^R D_{VF\mu}^R + L \leftrightarrow R) & f_{4b} &= \frac{1}{32e} (F_{Vf\mu}^R D_{\gamma Ff}^R D_{VF\mu}^L + L \leftrightarrow R) \\
f_{4c} &= \frac{1}{32e} (F_{Vf\mu}^R D_{\gamma Ff}^L D_{VF\mu}^L + L \leftrightarrow R) & f_{4d} &= \frac{1}{32e} (F_{Vf\mu}^R D_{\gamma Ff}^L D_{VF\mu}^R + L \leftrightarrow R) \\
f_{5a} &= -\frac{C_{VF'\mu}}{32eM_F^2} (D_{VF'\mu}^L D_{\gamma\mu^*\mu}^L + L \leftrightarrow R) & f_{5b} &= -\frac{C_{VF'\mu}}{32eM_F^2} (D_{VF'\mu}^L D_{\gamma\mu^*\mu}^R + L \leftrightarrow R) \\
f_{6a} &= -\frac{C_{VF'\mu^*}}{32eM_F^2} (D_{VF'\mu}^L D_{\gamma\mu^*\mu}^L + L \leftrightarrow R) & f_{6b} &= -\frac{C_{VF'\mu^*}}{32eM_F^2} (D_{VF'\mu}^R D_{\gamma\mu^*\mu}^L + L \leftrightarrow R) \\
f_{7a} &= -\frac{1}{32eM_F^2} (F_{Vf\mu}^R D_{V\mu^*f}^R D_{\gamma\mu^*\mu}^R + L \leftrightarrow R) & f_{7b} &= -\frac{1}{32eM_F^2} (F_{Vf\mu}^R D_{V\mu^*f}^R D_{\gamma\mu^*\mu}^L + L \leftrightarrow R) \\
f_{7c} &= -\frac{1}{32eM_F^2} (F_{Vf\mu}^R D_{V\mu^*f}^L D_{\gamma\mu^*\mu}^L + L \leftrightarrow R) & f_{7d} &= -\frac{1}{32eM_F^2} (F_{Vf\mu}^R D_{V\mu^*f}^L D_{\gamma\mu^*\mu}^R + L \leftrightarrow R) \\
f_{8a} &= -\frac{1}{32eM_F^2} (F_{Vf\mu}^L D_{V\mu^*f}^L D_{\gamma\mu^*\mu}^L + L \leftrightarrow R) & f_{8b} &= -\frac{1}{32eM_F^2} (F_{Vf\mu}^R D_{V\mu^*f}^L D_{\gamma\mu^*\mu}^L + L \leftrightarrow R) \\
f_{8c} &= -\frac{1}{32eM_F^2} (F_{Vf\mu}^R D_{V\mu^*f}^R D_{\gamma\mu^*\mu}^L + L \leftrightarrow R) & f_{8d} &= -\frac{1}{32eM_F^2} (F_{Vf\mu}^L D_{V\mu^*f}^R D_{\gamma\mu^*\mu}^L + L \leftrightarrow R)
\end{aligned}$$

References

- [1] H. N. Brown *et al.* [Muon g-2 Collaboration], Phys. Rev. Lett. **86**, 2227 (2001).
- [2] See for example, A. Dedes and H. E. Haber, hep-ph/0105014 and the references therein.
- [3] Some of the references are: J. Erler and M. Luo, Phys. Rev. Lett. **87**, 071804 (2001); A. Czarnecki and W. J. Marciano, Phys. Rev. D **64**, 013014 (2001); F. J. Yndurain, hep-ph/0102312; S. Narison, Phys. Lett. B **513**, 53 (2001); F. Jegerlehner, hep-ph/0104304; K. Melnikov, hep-ph/0105267.
- [4] For a review see, R. S. Chivukula, “Lectures on technicolor and compositeness,” hep-ph/0011264.
- [5] For recent works see for example: K. D. Lane, hep-ph/0102131; P. Das, S. Kumar Rai and S. Raychaudhuri, hep-ph/0102242; Z. H. Xiong and J. M. Yang, Phys. Lett. B **508**, 295 (2001); X. Calmet, H. Fritzsch and D. Holtmannspotter, Phys. Rev. D

- 64**, 037701 (2001); M. B. Einhorn and J. Wudka, Phys. Rev. Lett. **87**, 071805 (2001); C. x. Yue, Q. j. Xu and G. l. Liu, J. Phys. G **27**, 1807 (2001); Y. B. Dai, C. S. Huang and A. Zhang, hep-ph/0103317; X. Calmet, hep-ph/0108079.
- [6] P. Mery, S. E. Moubarik, M. Perrottet and F. M. Renard, Z. Phys. C **46**, 229 (1990).
 - [7] S. J. Brodsky and S. D. Drell, Phys. Rev. D **22** (1980) 2236; R. Escribano and E. Masso, Eur. Phys. J. C **4**, 139 (1998).
 - [8] J. H. Kuhn and P. M. Zerwas, Phys. Lett. B **147**, 189 (1984); J. H. Kuhn, H. D. Tholl and P. M. Zerwas, Phys. Lett. B **158**, 270 (1985); F. Cesi, ROME-432-1985.
 - [9] F. Boudjema, A. Djouadi and J. L. Kneur, Z. Phys. C **57**, 425 (1993).
 - [10] F. M. Renard, Phys. Lett. B **116**, 264 (1982).
 - [11] M. C. Gonzalez-Garcia and S. F. Novaes, Phys. Lett. B **389**, 707 (1996)
 - [12] A. Czarnecki and W. J. Marciano, hep-ph/0010194.
 - [13] B. Vachon, Nucl. Phys. Proc. Suppl. **98**, 148 (2001); D. E. Groom *et al.* [Particle Data Group Collaboration], Eur. Phys. J. C **15**, 1 (2000).
 - [14] E. Boos, A. Vologdin, D. Toback and J. Gaspard, hep-ph/0111034.
 - [15] O. J. Eboli, S. M. lietti and P. Mathews, hep-ph/0111001.